

Applicant : Paul G. Yock, et al.
Appl. No. : 10/776,037
Examiner : Marvich, Maria
Docket No. : 13854.4004

EXHIBIT A

37. (Amended) A method of locally administering an active agent to a host, said method comprising:

retroinfusing a fluid into a vascular vessel of said host under conditions sufficient for the fluid to produce a disruption in said vessel and infusing said agent into an interstitial space of said host through said disruption and locally administering said agent to said host through said disruption.

44. (Amended) A method of locally administering an active agent to a host, said method comprising:

retroinfusing a fluid into a vein of said host under conditions sufficient for the fluid to produce a disruption in said vein and infusing said agent into an interstitial space of said host through said disruption so that said agent is locally administered to said host.

51. (Amended) A method of locally administering an active agent to a host, said method comprising:

retroinfusing a fluid into a vein of said host with a catheter and at a pressure sufficient for the fluid to produce a disruption in said vein and infusing said agent into an interstitial space proximal to the vein through said disruption;

whereby said agent is locally administered to said host.

56. (Amended) A method of locally administering an active agent to a host, said method comprising:

Applicant : Paul G. Yock, et al.
Appl. No. : 10/776,037
Examiner : Marvich, Maria
Docket No. : 13854.4004

retroinfusing said agent into a vascular vessel of said host under conditions sufficient for the agent or a fluid delivery vehicle thereof to produce at least a mechanical stress on said vessel, which stress facilitates the transport of said agent through the wall of said vessel so that said agent is locally administered to said host,
wherein said method further comprises administration of energy to said vessel.

67. (Amended) A method of locally administering an active agent to a host, said method comprising:

retroinfusing said agent into a vascular vessel of said host under conditions sufficient for the agent or a fluid delivery vehicle thereof to at least distend said vessel, which distention facilitates the transport of said agent through the wall of said vessel so that said agent is locally administered to said host,
wherein said method further comprises administration of energy to said vessel.

78. (Amended) A method of locally administering an active agent to a host, said method comprising:

retroinfusing a fluid into a vascular vessel of said host under conditions sufficient for the agent or a fluid delivery vehicle thereof to produce a mechanical stress in said vessel, which stress facilitates the transport of said agent through a wall of said vessel so that said agent is locally administered to said host,
wherein said method further comprises administration of energy to said vessel.

Applicant : Paul G. Yock, et al.
Appl. No. : 10/776,037
Examiner : Marvich, Maria
Docket No. : 13854.4004

90. (Amended) A method of locally administering an active agent to a host, said method comprising:

retroinfusing a fluid into a vascular vessel of said host under conditions sufficient for the agent or a fluid delivery vehicle thereof to at least distend said vessel, which distention facilitates the transport of said agent through the wall of said vessel so that said agent is locally administered to said host

wherein said method further comprises administration of energy to said vessel.